

An inverse function theorem for Colombeau tame Frölicher–Kriegl maps

by

Seppo I. Hiltunen

Abstract. For $k = 1, 2, \dots, \infty$ and a Frölicher–Kriegl order k Lipschitz differentiable map $f : E \supseteq U \rightarrow E$ having derivative at $x_0 \in U$ a linear homeomorphism $E \rightarrow E$ and satisfying a Colombeau type tameness condition, we prove that x_0 has a neighbourhood $V \subseteq U$ with $f|_V$ a local order k Lipschitz diffeomorphism. As a corollary we obtain a similar result for Keller C_c^∞ maps with E in a class including Fréchet and Silva spaces. We also indicate a procedure for verifying the tameness condition for maps of the type $x \mapsto \varphi \circ [\text{id}, x]$ and spaces $E = C^\infty(Q)$ when Q is compact by considering the case $Q = [0, 1]$. Our considerations are motivated by the wish to try to retain something valuable in an interesting but defective treatment of integrability of Lie algebras by J. Leslie.

In [9; Theorem 4.1, p. 439] an interesting assertion is given which via [9; Lemma 4.2, p. 441] is based on [9; Theorem 2.2, p. 430] whose proof in turn is omitted for the most part with the exception of just few hints. It even seems that it cannot be proved unless one there requires the space E to be suitably restricted. To fill at least this gap in [9], we prove Theorem 8 below which then gives Corollary 9 as a replacement for [9; Theorem 2.2] when E is cm-convenient.

Proposition 10 below should serve as a prototype for assertions guaranteeing that Corollary 9 can be applied to maps of the type $x \mapsto \varphi \circ [\text{id}, x]$ of spaces $C^\infty(Q)$ when Q is compact. Below, we shall use the notational conventions of [3] from which we in particular recall the following

1 Conventions. Letting $\mathbf{R}$ be the topological field of real numbers, the class of all real locally convex Hausdorff topological vector spaces is $\text{LCS}(\mathbf{R})$. For $E \in \text{LCS}(\mathbf{R})$ we have $E = (X, \mathcal{T})$ where $X = \sigma_{r,d}E$ is the underlying “abstract” real vector space, and $\mathcal{T} = \tau_{r,d}E$ is the (locally convex) topology for the underlying set $v_s E$. The filter of $\mathcal{T}$ -neighborhoods of the zero vector $\mathbf{0}_E$ is $\mathcal{N}_o E$, and the von Neumann bornology, the set of all bounded sets is $\mathcal{B}_s E$.

A vector map of spaces in $\text{LCS}(\mathbf{R})$ is any triplet $\tilde{f} = (E, F, f)$ such that $E, F \in \text{LCS}(\mathbf{R})$ and f is a function with $f \subseteq (v_s E) \times (v_s F)$. The function value of f at x is $f \backslash x$, instead of the conventional notation “ $f(x)$ ”. The derivative at x of a Gateaux differentiable map $\tilde{f} = (E, F, f)$ is $\tilde{f}'(x)$.

By definition, we have $\text{dom}^2 \Gamma = \text{dom}(\text{dom } \Gamma)$ for any class Γ .

For $k \in \infty^+ = \mathbb{N}_o \cup \{\infty\}$, we below consider the differentiability classes $\mathcal{Lip}_{\text{FKt}}^k$ and $C_c^k(\mathbf{R})$. The latter one has as its members exactly the maps (E, F, f) with $E, F \in \text{LCS}(\mathbf{R})$ and $\text{dom } f \in \tau_{r,d}E$ such that for $l \in k + 1$, the order l variation

2000 *Mathematics Subject Classification.* 58C15, 46A61, 46T20.

Key words and phrases. Colombeau tame, Frölicher–Kriegl Lipschitz differentiable, bornological locally convex, locally complete, bornological generator, inverse function theorem.

$\delta_{EF}^l f$ has $\text{dom}(\delta_{EF}^l f) = (v_s E)^{l+1} \cap \{x : x \setminus \emptyset \in \text{dom} f\}$, and $\delta_{EF}^l f$ is continuous $\tau_{rd}(E^{l+1}]_{\text{tvs}}) \rightarrow \tau_{rd} F$. For more information, see [3; Section 3] and [7].

The class $\mathcal{Lip}_{\text{FKt}}^k$ has as its elements exactly the maps (E, F, f) such that the spaces $E, F \in \text{LCS}(\mathbf{R})$ are bornological and locally (i.e. Mackey) complete, see [6; p. 196] or [8; Lemma 2.2, p. 15], and $f : E \supseteq \text{dom} f \rightarrow F$ is $\mathcal{Lip}^k$ in the sense of [2; pp. 83, 99] or [8; Definition 12.1, p. 118]. Our choice corresponds to the one made in [2] where the spaces are bornological (locally convex) which is not required in [8]. This has the consequence that for any fixed $k \in \infty^+$ the class $\mathcal{Lip}_{\text{xzt}}^k \setminus \{(\mathbf{R}, E)\}$ of local k^{th} order Lipschitz differentiable curves in E uniquely determines E in the class $\text{LCS}(\mathbf{R}) \cap \{F : \sigma_{rd} E = \sigma_{rd} F \text{ and } \mathcal{B}_s E = \mathcal{B}_s F\}$ when $\text{xz} = \text{FK}$, but not when $\text{xz} = \text{KM}$.

For $E \in \text{LCS}(\mathbf{R})$, a set $U \subseteq v_s E$ we call *mopen* in E if and only if for all $x \in U$ and $B \in \mathcal{B}_s E$ there is $\delta \in \mathbb{R}^+$ with $tB \subseteq U - x$ for $0 \leq t \leq \delta$. The set $\tau_{\text{Mac}} E = \{U : U \text{ mopen in } E\}$ then is a topology for $v_s E$ with $\tau_{rd} E \subseteq \tau_{\text{Mac}} E$, equality here holding if E is metrizable, cf. [2; Remark 2.4.5, p. 52] where $\tau_{\text{Mac}} E$ is called the *Mackey closure topology*. For $(E, F, f) \in \mathcal{Lip}_{\text{FKt}}^k$ it holds that $\text{dom} f \in \tau_{\text{Mac}} E$ and that f is continuous $\tau_{\text{Mac}} E \rightarrow \tau_{\text{Mac}} F$, and further $(E, F, f|_U) \in \mathcal{Lip}_{\text{FKt}}^k$ for any $U \in \tau_{\text{Mac}} E$. For these facts, we refer the reader to see [2; Proposition 2.3.7, p. 44, Corollary 4.1.7, p. 85, Proposition 4.3.2, p. 99].

For short, a space $E \in \text{LCS}(\mathbf{R})$ we call *cm-convenient* iff also E is bornological and locally complete, and such that $\tau_{\text{Mac}}(E^k]_{\text{tvs}}) = \tau_{rd}(E^k]_{\text{tvs}})$ holds for $k \in \mathbb{N}_0$. By the above, all Fréchet spaces are cm-convenient, and by [5; Theorem 7.3.2(1), p. 99] also Silva spaces are cm-convenient, cf. [2; Theorem 6.1.4, p. 190].

2 Proposition. *Let $\tilde{f} = (E, F, f)$ where $F \in \text{dom}^2 \mathcal{Lip}_{\text{FKt}}^0$ and E is cm-convenient. The equivalence $\tilde{f} \in \mathcal{Lip}_{\text{FKt}}^\infty \Leftrightarrow \tilde{f} \in C_c^\infty(\mathbf{R})$ then holds.*

Proof. First letting $\tilde{f} \in \mathcal{Lip}_{\text{FKt}}^\infty$, for $k \in \mathbb{N}_0$ by [2; Theorem 4.3.27, p. 112], for the variation map $\delta^k \tilde{f} = (E^{k+1}]_{\text{tvs}}, F, \delta_{EF}^k f)$ we have $\delta^k \tilde{f} \in \mathcal{Lip}_{\text{FKt}}^0$ with $\text{dom}(\delta_{EF}^k f) = (v_s E)^{k+1} \cap \{x : x \setminus \emptyset \in \text{dom} f\}$. By [2; Corollary 4.1.7, p. 85], hence $\delta_{EF}^k f$ is continuous $\tau_{\text{Mac}}(E^{k+1}]_{\text{tvs}}) \rightarrow \tau_{\text{Mac}} F$, whence by $\tau_{rd} F \subseteq \tau_{\text{Mac}} F$, also continuous $\tau_{rd}(E^{k+1}]_{\text{tvs}}) \rightarrow \tau_{rd} F$. So we get $\delta^k \tilde{f} \in C_c^0(\mathbf{R})$. Here $k \in \mathbb{N}_0$ being arbitrary, we get $\tilde{f} \in C_c^\infty(\mathbf{R})$. Conversely, then letting $\tilde{f} \in C_c^\infty(\mathbf{R})$, for an arbitrary $c \in \mathcal{Lip}_{\text{FKt}}^\infty \setminus \{(\mathbf{R}, E)\} = C_c^\infty(\mathbf{R}) \setminus \{(\mathbf{R}, E)\}$ the chain rule gives the result that $f \circ c \in C_c^\infty(\mathbf{R}) \setminus \{(\mathbf{R}, F)\}$. Directly by definition, this further gives $\tilde{f} \in \mathcal{Lip}_{\text{FKt}}^\infty$, in view of [2; Lemma 4.3.1, p. 99]. $\square$

3 Definitions. For $E \in \text{LCS}(\mathbf{R})$, a set $\mathcal{B} \subseteq \mathcal{B}_s E \setminus \{\emptyset\}$ of absolutely convex closed sets we call a *bornological generator* for E iff for all $A, B \in \mathcal{B}$ there is $C \in \mathcal{B}$ with $A \cup B \subseteq C$, and every $A \in \mathcal{B}_s E$ has some $t \in \mathbb{R}^+$ and $B \in \mathcal{B}$ with $A \subseteq tB$.

For any real vector space X and any nonempty absolutely convex B in X and any $x \in v_s X$ with $S_{XB} x = \mathbb{R}^+ \cap \{t : t^{-1}x \in B\}$ we let $\|x\|_{XB} = \inf(S_{XB} x)$, hence having $\|x\|_{XB} = +\infty$ in case $x \notin S$ for the linear span $S = \tau_{rd} X[\mathbb{R}^+ \times B] = \{tv : t \in \mathbb{R}^+ \text{ and } v \in B\}$. In particular, if we have $X = \sigma_{rd} E$ with E locally complete and also B is $\tau_{rd} E$ -closed with $B \in \mathcal{B}_s E$, by [6; Proposition 10.2.1, p. 197] then $(X|_S, \langle \|x\|_{XB} : x \in S \rangle)$ is a normed Banach space with B its closed unit ball. The corresponding *Banachable* locally convex topological vector space is $X_B = (X|_S, \mathcal{T})$ for $\mathcal{T} = \mathcal{P}_s S \cap \{U : \forall x \in U ; \exists \varepsilon \in \mathbb{R}^+ ; \varepsilon B \subseteq U - x\}$. Note that we have $\tau_{\text{Mac}} E \cap S \subseteq \mathcal{T}$.

The standard argument in [10; Theorem 10.7, pp. 231–232] gives the following

4 Lemma. *Let $0 \leq \varepsilon < 1$, and let (X, ν) be a normed Banach space. If also ℓ is linear $X \rightarrow X$ with $\nu(\ell \backslash x - x) \leq \varepsilon(\nu \backslash x)$ for all $x \in \nu_s X$, then $\ell^{-\iota}$ is linear $X \rightarrow X$ with*

$$\nu \circ \ell^{-\iota} \backslash x \leq (1 - \varepsilon)^{-1}(\nu \backslash x) \quad \text{and}$$

$$\nu \backslash (\ell^{-\iota} \backslash x - x) \leq (1 - \varepsilon)^{-1} \varepsilon(\nu \backslash x) \quad \text{for all } x \in \nu_s X.$$

5 Definitions. We let $\text{Colo}_A(\tilde{f}, y_0, \varepsilon, \mathcal{B})$ mean that there are E, f such that $\tilde{f} = (E, E, f)$ with $\tilde{f}$ a Gateaux differentiable map and $E \in \text{LCS}(\mathbf{R})$ bornological and locally complete and $\varepsilon \in \mathbb{R}^+$ and $\mathcal{B}$ a bornological generator for E such that for $v \in B \in \mathcal{B}$ and for $y_1 = y_0 + 2v$ we have $y_1 \in (\text{dom } f) \cap (v_s(\sigma_{rd} E_B))$, and also the inclusion $(\tilde{f}'(y_1) - \tilde{f}'(y_0)) \backslash B \subseteq \varepsilon \tilde{f}'(y_0) \backslash B$ holds.

Let $\text{Colo}_0(\tilde{f}, y_0, \varepsilon, \mathcal{B})$ mean that $\text{Colo}_A(\tilde{f}, y_0, \varepsilon, \mathcal{B})$ holds and $\tilde{f}'(y_0)$ is a linear homeomorphism $E \rightarrow E$. For $\tilde{f} = (E, E, f)$, letting $\tilde{f}|_{\text{map}} V = (E, E, f|_V)$, a vector map $\tilde{f}$ we say to be *Colombeau₀-tame* at y_0 iff for all V with $y_0 \in V \in \tau_{rd} E$ there is some $\mathcal{B}$ such that $\text{Colo}_0(\tilde{f}|_{\text{map}} V, y_0, \frac{1}{2}, \mathcal{B})$ holds.

6 Remark. We note some facts to be used below.

(a) Assuming $\text{Colo}_A(\tilde{f}, y_0, \varepsilon, \mathcal{B})$ to hold with $\tilde{f} = (E, E, f)$ and $X = \sigma_{rd} E$ and $B \in \mathcal{B}$, since $\mathbf{0}_E \in B$ by circledness of B following from its absolute convexity, we have $y_0 = y_0 + 2\mathbf{0}_E \in \text{dom } f \cap v_s(X_B)$, and consequently $Q \subseteq v_s(X_B)$ holds for the closed convex set $Q = y_0 + 2B$. Further $(\tau_{rd}(X_B), \tau_{rd} E, f|_Q)$ is a *topological map*, i.e. we have $f|_Q$ continuous $\tau_{rd}(X_B) \rightarrow \tau_{rd} E$.

To see this, arbitrarily fixing $y_1 \in Q$ and a closed convex $V \in \mathcal{N}_o E$, there should be some $N \in \mathcal{N}_{bh}(y_1, \tau_{rd}(X_B) \cap Q)$ with $f \backslash N \subseteq f \backslash y_1 + V$. For this, we first note that $\tilde{f}'(y_0)$ being a continuous linear map $E \rightarrow E$, it is bornological $\mathcal{B}_s E \rightarrow \mathcal{B}_s E$, and hence we have $\tilde{f}'(y_0) \backslash B \in \mathcal{B}_s E$, whence further there is some $\varepsilon_1 \in \mathbb{R}^+$ with $\varepsilon_1 \tilde{f}'(y_0) \backslash B \subseteq V$. With $\lambda = (1 + \varepsilon)^{-1} \varepsilon_1$, now taking $N = (y_1 + \lambda B) \cap Q$, for $y_2 = y_1 + v \in N$ we see $f \backslash y_2 \in f \backslash y_1 + V$ to hold as follows.

For $I = [0, 1]$ and $c = \langle f \backslash (y_1 + tv) - f \backslash y_1 : t \in I \rangle$, we have c a differentiable curve in E with $c \backslash 0 = \mathbf{0}_E$ and $c \backslash 1 = f \backslash y_2 - f \backslash y_1$. By the mean value theorem, it hence suffices that $\text{rng}(D_E c) \subseteq V$. This is the case since for $t \in I$ we have

$$\begin{aligned} D_E c \backslash t &= \tilde{f}'(y_1 + tv) \backslash v \in (\tilde{f}'(y_1 + tv) - \tilde{f}'(y_0)) \backslash (\lambda B) + \tilde{f}'(y_0) \backslash (\lambda B) \\ &\subseteq \lambda(\varepsilon \tilde{f}'(y_0) \backslash B + \tilde{f}'(y_0) \backslash B) \subseteq \lambda(1 + \varepsilon) \tilde{f}'(y_0) \backslash B \subseteq \varepsilon_1 \tilde{f}'(y_0) \backslash B \subseteq V. \end{aligned}$$

(b) Assuming $\text{Colo}_0(\tilde{f}, y_0, \varepsilon, \mathcal{B})$ to hold with $\tilde{f} = (E, E, f)$ and $X = \sigma_{rd} E$ and $\varepsilon < 1$, for $y_1 \in y_0 + 2 \bigcup \mathcal{B}$ we have $\tilde{f}'(y_1)$ a *linear homeomorphism* $E \rightarrow E$. For this, since we assume E to be bornological, it suffices that $\tilde{f}'(y_1)$ is bijective $v_s E \rightarrow v_s E$, and that $\tilde{f}'(y_1)$ and $(\tilde{f}'(y_1))^{-\iota}$ are bornological $\mathcal{B}_s E \rightarrow \mathcal{B}_s E$.

First, to show indirectly that $\tilde{f}'(y_1)$ is injective, if not, there is $v \in v_s E \setminus \{\mathbf{0}_E\}$ with $\tilde{f}'(y_1) \backslash v = \mathbf{0}_E$. There further is $B_1 \in \mathcal{B}$ with $y_1 \in y_0 + 2B_1$, and there is $B_2 \in \mathcal{B}$ with $v \in v_s(X_{B_2})$. We then find $B_3 \in \mathcal{B}$ with $B_1 \cup B_2 \subseteq B_3$, and we have $(\tilde{f}'(y_1) - \tilde{f}'(y_0)) \backslash B_3 \subseteq \varepsilon \tilde{f}'(y_0) \backslash B_3$. For $\ell = (\tilde{f}'(y_0))^{-\iota} \circ (\tilde{f}'(y_1))|_{v_s(X_{B_3})}$ it follows that $0 < \|v\|_{X_{B_3}} = \|\ell \backslash v - v\|_{X_{B_3}} \leq \varepsilon \|v\|_{X_{B_3}} < \|v\|_{X_{B_3}}$.

To get $v_s E \subseteq \text{rng}(\tilde{f}'(y_1))$, given $v \in v_s E$, deducing as above, we find B_3 with $(\tilde{f}'(y_1) - \tilde{f}'(y_0)) \backslash B_3 \subseteq \varepsilon \tilde{f}'(y_0) \backslash B_3$, and now $\tilde{f}'(y_0)^{-\iota} \backslash v \in v_s(X_{B_2})$. For $x \in v_s(X_{B_3})$ then $\|\ell \backslash x - x\|_{X_{B_3}} \leq \varepsilon \|x\|_{X_{B_3}}$ whence Lemma 4 gives $\tilde{f}'(y_0)^{-\iota} \backslash v \in v_s(X_{B_3}) \subseteq \text{rng } \ell$, and so there is $u \in v_s(X_{B_3}) \subseteq v_s E$ with $\tilde{f}'(y_1) \backslash u = v$.

To prove that $\tilde{f}'(y_1)$ is bornological $\mathcal{B}_s E \rightarrow \mathcal{B}_s E$, given $B \in \mathcal{B}_s E$, as above, we find B_3 with now also having $B \subseteq \lambda B_3$ for a suitable $\lambda \in \mathbb{R}^+$. Then we obtain $\tilde{f}'(y_1) \setminus B \subseteq \lambda \tilde{f}'(y_1) \setminus B_3 \subseteq \lambda(1 + \varepsilon) \tilde{f}'(y_0) \setminus B_3 \in \mathcal{B}_s E$. To get the assertion for $(\tilde{f}'(y_1))^{-\iota}$, we arrange $B \subseteq \lambda \tilde{f}'(y_0) \setminus B_3$, and by Lemma 4 we obtain $\ell^{-\iota} \setminus B_3 \subseteq (1 - \varepsilon)^{-1} B_3$, whence $\tilde{f}'(y_1)^{-\iota} \setminus B \subseteq \lambda(1 - \varepsilon)^{-1} B_3 \in \mathcal{B}_s E$ follows.

(c) Assuming $\text{Colo}_0(\tilde{f}, y_0, \varepsilon, \mathcal{B})$ to hold with $\tilde{f} = (E, E, f)$ and $X = \sigma_{rd} E$ and $\ell = (\tilde{f}'(y_0))^{-\iota}$, for $f_1 = \langle y - \ell \circ f \setminus y : y \in \text{dom } f \rangle$, and for $y_8, y_9 \in y_0 + 2B$ with $B \in \mathcal{B}$, we have $\|f_1 y_8 - f_1 y_9\|_{XB} \leq \varepsilon \|y_8 - y_9\|_{XB}$. Indeed, putting $v = y_8 - y_9$ and $I = [0, 1]$, and considering in the space E the differentiable curve $c = \langle tv - \ell \setminus (f \setminus (y_9 + tv) - f \setminus y_9) : t \in I \rangle$, for which we have $c \setminus 0 = \mathbf{0}_E$ and $c \setminus 1 = f_1 y_8 - f_1 y_9$ and $D_E c = \langle v - \ell \circ (\tilde{f}'(y_9 + tv)) \setminus v : t \in I \rangle$, to get the assertion, by the mean value theorem, for arbitrarily fixed $t \in I$, it suffices that $\|D_E c \setminus t\|_{XB} \leq \varepsilon \|v\|_{XB}$. This is the case, since for $y_7 = y_9 + tv$ and for $\ell_1 = \ell \circ (\tilde{f}'(y_7))$, we have the inclusion $(\tilde{f}'(y_7) - \tilde{f}'(y_0)) \setminus B \subseteq \varepsilon \tilde{f}'(y_0) \setminus B$, which further gives $\|D_E c \setminus t\|_{XB} = \|\ell_1 \setminus v - v\|_{XB} \leq \varepsilon \|v\|_{XB}$.

7 Lemma. Let $\text{Colo}_0(\tilde{f}, y_0, \varepsilon, \mathcal{B})$ hold with $0 \leq \varepsilon \leq \frac{1}{2}$ and $\tilde{f} = (E, E, f)$ and $X = \sigma_{rd} E$ and $x_0 = f \setminus y_0$. Also let $\ell = (\tilde{f}'(y_0))^{-\iota}$ and $Q = x_0 + \ell^{-\iota} \cup \mathcal{B}$. Then there is a function $g \subseteq f^{-\iota}$ with $x_0 \in \text{dom } g \in \tau_{rd} E \cap \mathcal{P}_s Q$, and in addition for every $B \in \mathcal{B}$ and for all $x_1, x_2 \in (x_0 + \ell^{-\iota} \setminus B) \cap \text{dom } g$ it holds that

$$\|g \setminus x_1 - g \setminus x_2\|_{XB} \leq (1 - \varepsilon)^{-1} \|\ell \setminus (x_1 - x_2)\|_{XB}.$$

Proof. Let $h = \{(x, y, z) : \exists x_1; (y, x_1) \in f \text{ and } \ell \setminus x - \ell \setminus x_1 + y = z\}$.

Then h is a function, and we further put $g_1 = \text{dom}^2 \Gamma$, where Γ is the set of all $(x, y, B, \mathbf{y})$ such that $B \in \mathcal{B}$ and $\mathbf{y} \in (v_s E)^{N_0}$ with $x \in x_0 + \tilde{f}'(y_0) \setminus B$ and $\mathbf{y} \setminus \emptyset = y_0$ and $\mathbf{y} \rightarrow y$ in $\text{top } \tau_{rd} E$ and $(x, \mathbf{y} \setminus i, \mathbf{y} \setminus i^+) \in h$ for all $i \in N_0$.

We note that g_1 is a function, since if $(x, y_i) \in g_1$ for $i=1,2$, we get $y_1 = y_2$ as follows. There are B_i and $\mathbf{y}_i$ with $(x, y_i, B_i, \mathbf{y}_i) \in \Gamma$. As we have $\mathbf{y}_i \setminus \emptyset = y_0$ and $(x, \mathbf{y}_i \setminus i, \mathbf{y}_i \setminus i^+) \in h$ for all $i \in N_0$, since h is a function, by induction we get $\mathbf{y}_1 \setminus i = \mathbf{y}_2 \setminus i$ for $i \in N_0$, and hence $\mathbf{y}_1 = \mathbf{y}_2$. Since $\tau_{rd} E$ is a Hausdorff topology, and as we have $\mathbf{y}_1 \rightarrow y_1$ in $\text{top } \tau_{rd} E$, it follows that $y_1 = y_2$.

We next prove that $\text{dom } g_1 = Q$. Trivially having $\text{dom } g_1 \subseteq Q$, arbitrarily given $x \in x_0 + \tilde{f}'(y_0) \setminus B$ with $B \in \mathcal{B}$, it suffices to show that there are $y, \mathbf{y}$ with $(x, y, B, \mathbf{y}) \in \Gamma$. To establish this, we construct $\mathbf{y}$ by the following recursion: fixing any $z_0 \in \mathbf{U} \setminus v_s E$, we require that $\mathbf{y} \setminus \emptyset = y_0$ and $\mathbf{y} \setminus i^+ = h \setminus (x, \mathbf{y} \setminus i)$ in case $(x, \mathbf{y} \setminus i) \in \text{dom } h$, otherwise putting $\mathbf{y} \setminus i^+ = z_0$, for all $i \in N_0$. Letting $(l)_A$ mean that $\|\mathbf{y} \setminus i^{++} - \mathbf{y} \setminus i^+\|_{XB} \leq \varepsilon \|\mathbf{y} \setminus i^+ - \mathbf{y} \setminus i\|_{XB} < +\infty$ holds for all $i \in l^+$, we first establish $\forall l \in N_0; (l)_A$ by induction as follows.

To get $(\emptyset)_A$, first note that $\mathbf{y} \setminus 1. = \mathbf{y} \setminus \emptyset^+ = \ell \setminus x - \ell \setminus x_0 + y_0$. To get $(x, \mathbf{y} \setminus 1.) \in \text{dom } h$, by our arrangements and Definitions 5 it suffices for $v_0 = \mathbf{y} \setminus 1. - y_0$ that $v_0 \in 2B$. This holds by $v_0 = \ell \setminus (x - x_0) \in \ell \setminus (\tilde{f}'(y_0) \setminus B) = B \subseteq 2B$, which also gives $\|v_0\|_{XB} \leq 1$. We hence have

$$\begin{aligned} \mathbf{y} \setminus 2. &= \mathbf{y} \setminus \emptyset^{++} = h \setminus (x, \mathbf{y} \setminus 1.) = \ell \setminus x - \ell \circ f \setminus (\mathbf{y} \setminus 1.) + \mathbf{y} \setminus 1., & \text{whence} \\ \mathbf{y} \setminus 2. - \mathbf{y} \setminus 1. &= \ell \setminus x - \ell \circ f \setminus (\mathbf{y} \setminus 1.) = \ell \setminus x_0 + \mathbf{y} \setminus 1. - y_0 - \ell \circ f \setminus (\mathbf{y} \setminus 1.) \\ &= v_0 - \ell \setminus (f \setminus (\mathbf{y} \setminus 1.) - f \setminus y_0) = f_1 y_8 - f_1 y_0, \end{aligned}$$

taking $y_8 = \mathbf{y} \setminus 1.$. Remark 6(c) now gives the assertion.

With $l \in N_0$ now assuming that $(l)_A$ holds, we prove $(l^+)_A$ as follows. Since we have $(l)_A$, for $v_1 = \mathbf{y} \setminus l^{++} - \mathbf{y} \setminus l^+$ and $v_2 = \mathbf{y} \setminus l^{+++} - \mathbf{y} \setminus l^{++}$ we only have to establish $\|v_2\|_{XB} \leq \varepsilon \|v_1\|_{XB}$. By $(l)_A$ we have

$$\| \mathbf{y}^{\setminus l^{++}} - y_0 \|_{XB} \leq \sum_{i \in l^{++}} \| \mathbf{y}^{\setminus i^+} - \mathbf{y}^{\setminus i} \|_{XB}$$

$$\leq \sum_{i \in l^{++}} \varepsilon^i \| v_0 \|_{XB} \leq \sum_{i \in l^{++}} \varepsilon^i \leq 2,$$
 (e) and consequently $(x, \mathbf{y}^{\setminus l^{++}}) \in \text{dom } h$, whence further

$$\mathbf{y}^{\setminus l^{+++}} = \ell^{\setminus} x - \ell \circ f^{\setminus}(\mathbf{y}^{\setminus l^{++}}) + \mathbf{y}^{\setminus l^{++}}.$$
 Also having

$$\mathbf{y}^{\setminus l^{++}} = \ell^{\setminus} x - \ell \circ f^{\setminus}(\mathbf{y}^{\setminus l^+}) + \mathbf{y}^{\setminus l^+},$$
 we get

$$v_2 = v_1 - \ell^{\setminus}(f^{\setminus}(\mathbf{y}^{\setminus l^{++}}) - f^{\setminus}(\mathbf{y}^{\setminus l^+})) = f_1^{\setminus}(\mathbf{y}^{\setminus l^{++}}) - f_1^{\setminus}(\mathbf{y}^{\setminus l^+}),$$
 whence again Remark 6(c) gives the assertion.

Now having obtained $\forall l \in \mathbb{N}_0; (l)_A$, we know that $\mathbf{y}$ is a Cauchy sequence in X_B , which is Banachable, hence complete by the assumption that E is locally complete. Hence, there is y with $\mathbf{y} \rightarrow y$ in $\text{top } \tau_{rd}(X_B)$, and consequently also $\mathbf{y} \rightarrow y$ in $\text{top } \tau_{rd}E$. We have now concluded the proof of $\text{dom } g_1 = Q$.

From the assumption that $\mathcal{B}$ is a bornological generator for E it follows that $\bigcup \mathcal{B}$ is an absolutely convex bornivore in E , hence also $\tilde{f}'(y_0)^{\setminus} \bigcup \mathcal{B}$ since $\tilde{f}'(y_0)$ is assumed to be a linear homeomorphism $E \rightarrow E$. Since E is assumed to be bornological, we have $\tilde{f}'(y_0)^{\setminus} \bigcup \mathcal{B} \in \mathcal{N}_o E$, whence it follows existence of some $U \in \tau_{rd}E$ with $x_0 \in U \subseteq Q$. Then putting $g = g_1|U$, we have g a function with $x_0 \in \text{dom } g = U \in \tau_{rd}E \cap \mathcal{P}_s Q$, recalling that $\mathcal{P}_s Q = \{ S : S \subseteq Q \}$.

We now proceed to prove $g \subseteq f^{\setminus \iota}$. For this considering arbitrary $(x, y) \in g$, we have $(x, y) \in g_1$, whence there are $B, \mathbf{y}$ with $(x, y, B, \mathbf{y}) \in \Gamma$. A slight rearrangement of the arguments used to establish $\text{dom } g_1 = Q$ shows that (e) holds. This gives $\mathbf{y} \in (y_0 + 2B)^{\mathbb{N}_0}$ whence by closedness of B we get $y \in y_0 + 2B \subseteq \text{dom } f$. Since for $P = (\tau_{rd}(X_B), \tau_{rd}E)$ we have $(P, f|(y_0 + 2B))$ a continuous map, also $(P, \ell \circ f|(y_0 + 2B))$ is such. For $i \in \mathbb{N}_0$ having $\mathbf{y}^{\setminus i^+} = \ell^{\setminus} x - \ell \circ f^{\setminus}(\mathbf{y}^{\setminus i}) + \mathbf{y}^{\setminus i}$, we get $y = \ell^{\setminus} x - \ell \circ f^{\setminus} y + y$, consequently $x = f^{\setminus} y$, and hence $(x, y) \in f^{\setminus \iota}$.

Fixing $B \in \mathcal{B}$ and $x_1, x_2 \in (x_0 + \ell^{\setminus \iota} B) \cap U$, for $y_i = g^{\setminus} x_i$ and $u = x_1 - x_2$ and $v = y_1 - y_2$ it remains to establish $\|v\|_{XB} \leq (1 - \varepsilon)^{-1} \|\ell^{\setminus} u\|_{XB}$. We have $y_i = \ell^{\setminus} x_i - \ell \circ f^{\setminus} y_i + y_i$, whence we get

$$v - \ell^{\setminus} u = v - \ell^{\setminus}(f^{\setminus} y_1 - f^{\setminus} y_2) = f_1^{\setminus} y_1 - f_1^{\setminus} y_2.$$

Noting that by the preceding paragraph we have $y_1, y_2 \in y_0 + 2B$, by Remark 6(c) we get

$$\|v - \ell^{\setminus} u\|_{XB} = \|f_1^{\setminus} y_1 - f_1^{\setminus} y_2\|_{XB} \leq \varepsilon \|v\|_{XB}$$

$$\text{and further } \|v\|_{XB} = \|v - \ell^{\setminus} u + \ell^{\setminus} u\|_{XB} \\ \leq \|v - \ell^{\setminus} u\|_{XB} + \|\ell^{\setminus} u\|_{XB} \leq \varepsilon \|v\|_{XB} + \|\ell^{\setminus} u\|_{XB},$$

whence finally $\|v\|_{XB} \leq (1 - \varepsilon)^{-1} \|\ell^{\setminus} u\|_{XB}$. $\square$

8 Theorem. *If $k \neq \emptyset$ and $\tilde{f} = (E, E, f) \in \mathcal{Lip}_{\text{FKt}}^k$ and $\tilde{f}$ is Colombeau₀-tame at y_0 , there is U with $y_0 \in U \in \tau_{\text{Mac}} E$ and $(E, E, (f|U)^{\setminus \iota}) \in \mathcal{Lip}_{\text{FKt}}^k$.*

Proof. Assuming the premise, let $x_0 = f^{\setminus} y_0$ and $X = \sigma_{rd} E$. Now, there is $\mathcal{B}_0$ such that $\text{Colo}_0(\tilde{f}|_{\text{map}} v_s E, y_0, \frac{1}{2}, \mathcal{B}_0)$ holds. Putting $Q_0 = y_0 + 2 \bigcup \mathcal{B}_0$, we show indirectly that $f|Q_0$ is injective. Indeed, if this does not hold, there are distinct $y_1, y_2 \in \text{dom } f \cap Q_0$ with $f^{\setminus} y_1 = f^{\setminus} y_2$, and we find some $B \in \mathcal{B}_0$ with $y_1, y_2 \in y_0 + 2B$. Letting $v = y_1 - y_2$, by Remark 6(c) we then get $\|v\|_{XB} = \|y_1 - y_2\|_{XB} = \|f_1^{\setminus} y_1 - f_1^{\setminus} y_2\|_{XB} \leq \frac{1}{2} \|y_1 - y_2\|_{XB} = \frac{1}{2} \|v\|_{XB}$ whence finally $0 < \|v\|_{XB} = 2 \|v\|_{XB} - \|v\|_{XB} \leq \|v\|_{XB} - \|v\|_{XB} = 0$, a contradiction.

Next, since $\bigcup \mathcal{B}_0$ is an absolutely convex bornivore in E , there is $V \in \tau_{rd} E$ with $y_0 \in V \subseteq Q_0$, whence there further is some $\mathcal{B}$ with $\text{Colo}_0(\tilde{f}|_{\text{map}} V, y_0, \frac{1}{2}, \mathcal{B})$. Letting g be as given by Lemma 7 above, and taking $U = \text{rng } g$, since $f|V$ is injective and $g \subseteq (f|V)^{\setminus \iota}$, and as $\text{dom } g \in \tau_{rd} E \subseteq \tau_{\text{Mac}} E$, in view of continuity

of $(\tau_{\text{Mac}} E, \tau_{\text{Mac}} E, f | V)$ it follows that $U = (f | V)^{-\iota} [\text{dom } g] \in \tau_{\text{Mac}} E$. Trivially having $y_0 \in U$, for $\tilde{g} = (E, E, g)$ it remains to establish $\tilde{g} \in \mathcal{Lip}_{\text{FKt}}^k$.

For this using [2; Theorem 4.8.4, p. 152], in view of Remark 6(b) it suffices that $\tilde{g} \in \mathcal{Lip}_{\text{FKt}}^0$. That is, for arbitrarily given $c \in \mathcal{Lip}_{\text{FKt}}^0 \{(\mathbf{R}, E)\}$ and $\gamma = g \circ c$, we should have $\gamma \in \mathcal{Lip}_{\text{FKt}}^0 \{(\mathbf{R}, E)\}$. To get this, for arbitrarily given $t_0 \in \text{dom } \gamma$ it suffices to show existence of $\delta \in \mathbb{R}^+$ and $B \in \mathcal{B}$ such that for $J =]t_0 - \delta, t_0 + \delta[$ we have $\gamma | J \in \mathcal{Lip}_{\text{FKt}}^0 \{(\mathbf{R}, X_B)\}$. To get this, we put $\mathcal{B}_1 = \ell^{-\iota} \mathcal{B}$, and first note that $c \backslash t_0 - y_0 \in \text{dom } g - y_0 \in \tau_{rd} E$. Hence, there is a real $r > 1$ with $r(c \backslash t_0 - y_0) \in \text{dom } g - y_0 \subseteq \bigcup \mathcal{B}_1$, whence further there is $B_2 \in \mathcal{B}_1$ with $r(c \backslash t_0 - y_0) \in B_2$. By [8; Corollary 1.8, p. 13] we then find $\delta \in \mathbb{R}^+$ and $B_1 \in \mathcal{B}_1$ with $B_2 \subseteq B_1$ and $c | J \in \mathcal{Lip}_{\text{FKt}}^0 \{(\mathbf{R}, X_{B_1})\}$. Since now $c \backslash t_0 - y_0 \in \text{Int}_{\tau_{rd}(X_{B_1})} B_1$, we may take δ smaller so that also $c \backslash J \subseteq y_0 + B_1$. For $B = \ell \backslash B_1$, the “in addition” part of Lemma 7 now gives $\gamma | J \in \mathcal{Lip}_{\text{FKt}}^0 \{(\mathbf{R}, X_B)\}$. $\square$

9 Corollary. *If $\tilde{f} = (E, E, f) \in C_c^\infty(\mathbf{R})$ with E a cm-convenient space, and if also $\tilde{f}$ is Colombeau₀-tame at x_0 , there is U with $x_0 \in U \in \tau_{rd} E$ and $(E, E, (f | U)^{-\iota}) \in C_c^\infty(\mathbf{R})$.*

Proof. Since $\tau_{\text{Mac}} E = \tau_{rd} E$ holds for cm-convenient E , the assertion immediately follows from Proposition 2 and Theorem 8 above. $\square$

To indicate the basic idea for proving Colombeau₀-tameness for maps of the type $x \mapsto \varphi \circ [\text{id}, x]$, and also to show that Colombeau₀-tameness despite of its strength is not too restrictive, we establish the following

10 Proposition. *Let $I = [0, 1]$ and $E = C^\infty(I)$, and also let $\varphi : I \times \mathbb{R} \rightarrow \mathbb{R}$ be smooth with $0 \notin \text{rng } \partial_2 \varphi$. For $f = \langle \varphi \circ [\text{id}, x] : x \in v_s E \rangle$ and for $\tilde{f} = (E, E, f)$, then $\tilde{f}$ is Colombeau₀-tame at every $x \in v_s E$.*

Proof. Assuming that $x \in V \in \tau_{rd} E$, there is $l_0 \in \mathbb{N}_0$ with the property that $x + V_0 \subseteq V$ for the set $V_0 = v_s E \cap \{z : \forall i \in l_0^+, s \in I; l_0 | z^{(i)} \backslash s| \leq 2\}$. For

$$\chi = I \times \mathbb{R} \times \mathbb{R} \cap \left\{ (s, \eta, t) : \begin{aligned} t &= 4(\partial_2 \varphi \backslash (s, x \backslash s))^{-1} \int_0^1 \partial_2^2 \varphi \backslash (s, x \backslash s + 2s_1 \eta) \, ds_1 \end{aligned} \right\},$$

we observe that $2(\tilde{f}'(x))^{-\iota} \circ (\tilde{f}'(x + 2u) - \tilde{f}'(x)) \backslash v = \chi \circ [\text{id}, u] \cdot u \cdot v$ for any $u, v \in v_s E$. To prove that $\tilde{f}$ is Colombeau₀-tame at x , it hence suffices to establish a bornological generator $\mathcal{B}$ for E such that we have $x \in v_s(\sigma_{rd} E_B)$ and

$$(*) \quad \chi \circ [\text{id}, u] \cdot u \cdot v \in B \subseteq \frac{1}{2} V_0 \quad \text{whenever } u, v \in B \in \mathcal{B}.$$

To get this, we make the following preparations and observations. Write

$$\mathbf{B} \mathbf{m} = v_s E \cap \{x : \forall i \in \mathbb{N}_0, s \in I; |x^{(i)} \backslash s| \leq \mathbf{m} \backslash i\},$$

where we generally require $\mathbf{m} \in (\mathbb{R}^+)^{\mathbb{N}_0}$ to be nondecreasing. For $i \in \mathbb{N}_0$, let $\text{Pi} = \mathbb{N}_0^{\times 2} \cap \{(i_1, i_2) : i_1 + i_2 \in i^+\}$, and for $\chi_1 \in v_s C^\infty(I \times \mathbb{R})$ and $u \in v_s E$ define the “jet” functions

$$J_2^i \chi_1 : I \times \mathbb{R} \ni \zeta \mapsto \{(i_1, i_2, \partial_1^{i_1} \partial_2^{i_2} \chi \backslash \zeta) : (i_1, i_2) \in \text{Pi}\} \in \mathbb{R}^{\text{Pi}}$$

$$\text{and } J_1^i u : I \ni s \mapsto \langle u^{(l)} \backslash s : l \in i^+ \rangle \in \mathbb{R}^{i+1}.$$

If we have a polynomial function $p : \mathbb{R}^{\text{Pi}} \times \mathbb{R}^i \times \mathbb{R}^i \rightarrow \mathbb{R}$ with the properties that for fixed ξ the map $(\eta, \zeta) \mapsto p \backslash (\eta, \zeta, \xi)$ is bilinear $\mathbb{R}^{\text{Pi}} \times_{\text{vs}} \times_{\text{vs}} (\mathbb{R}^i \times_{\text{vs}}) \rightarrow \mathbb{R}$, and for all $\chi_1 \in v_s C^\infty(I \times \mathbb{R})$ and $u, v \in v_s E$ we have

$$(s) \quad \begin{aligned} (\chi_1 \circ [\text{id}, u] \cdot u \cdot v)^{(i)} &= \partial_2 \chi_1 \circ [\text{id}, u] \cdot u^{(i)} \cdot u \cdot v + \chi_1 \circ [\text{id}, u] \cdot u^{(i)} \cdot v \\ &+ \chi_1 \circ [\text{id}, u] \cdot u \cdot v^{(i)} + p \circ [J_2^i \chi_1 \circ [\text{id}, u], J_1^{i-1} v, J_1^{i-1} u], \end{aligned}$$

$$\begin{aligned}
 \text{it follows that } & (\chi_1 \circ [\text{id}, u] \cdot u \cdot v)^{(i+1)} = \\
 & \partial_1 \partial_2 \chi_1 \circ [\text{id}, u] \cdot u^{(i)} \cdot u \cdot v + \partial_2^2 \chi_1 \circ [\text{id}, u] \cdot u' \cdot u^{(i)} \cdot u \cdot v \\
 & + \partial_2 \chi_1 \circ [\text{id}, u] \cdot u^{(i+1)} \cdot u \cdot v + \partial_2 \chi_1 \circ [\text{id}, u] \cdot u^{(i)} \cdot u' \cdot v \\
 & + \partial_2 \chi_1 \circ [\text{id}, u] \cdot u^{(i)} \cdot u \cdot v' + \partial_1 \chi_1 \circ [\text{id}, u] \cdot u^{(i)} \cdot v \\
 & + \partial_2 \chi_1 \circ [\text{id}, u] \cdot u' \cdot u^{(i)} \cdot v + \chi_1 \circ [\text{id}, u] \cdot u^{(i+1)} \cdot v \\
 & + \chi_1 \circ [\text{id}, u] \cdot u^{(i)} \cdot v' + \partial_1 \chi_1 \circ [\text{id}, u] \cdot u \cdot v^{(i)} \\
 & + \partial_2 \chi_1 \circ [\text{id}, u] \cdot u' \cdot u \cdot v^{(i)} + \chi_1 \circ [\text{id}, u] \cdot u' \cdot v^{(i)} \\
 & + \chi_1 \circ [\text{id}, u] \cdot u \cdot v^{(i+1)} + p \circ [J_2^i \partial_1 \chi_1 \circ [\text{id}, u], J_1^{i-1} \cdot v, J_1^{i-1} \cdot u] \\
 & + p \circ [J_2^i \partial_2 \chi_1 \circ [\text{id}, u], J_1^{i-1} \cdot v, J_1^{i-1} \cdot u] \cdot u' \\
 & + p \circ [J_2^i \chi_1 \circ [\text{id}, u], J_1^{i-1} \cdot v', J_1^{i-1} \cdot u] \\
 & + \partial_3 p \circ [J_2^i \chi_1 \circ [\text{id}, u], J_1^{i-1} \cdot v, J_1^{i-1} \cdot u] \cdot (J_1^{i-1} \cdot u').
 \end{aligned}$$

Omitting the details, from the preceding one sees that by a suitable recursion one can construct $\mathbf{P} \in \mathbf{U}^N$ such that for $(i, p) \in \mathbf{P}$ we have $p = \mathbf{P} \cdot i$ a polynomial as above such that (s) holds for the appropriate χ_1, u, v . Letting $\mathbf{R}is$ denote

$$\begin{aligned}
 & \sup \{ |\mathbf{P} \cdot i^+ \cdot (\xi, \eta, \zeta)| : i : \xi \in J_2^{i+1} \cdot \chi[I^{\times 2}] \} \\
 & \text{and } \sup \{ |r| : r \in \text{rng}(\eta \cup \xi) \} \leq s \},
 \end{aligned}$$

and constructing $\rho \in \mathbf{U}^{N_0}$ by the recursion

$$\begin{aligned}
 \rho \cdot \emptyset &= \langle \max \{s, \mathbf{R} \emptyset s\} : s \in \mathbb{R}_+ \rangle \quad \text{and} \\
 \rho \cdot i^+ &= \langle \max \{ \rho \cdot i \cdot s, \mathbf{R} i^+ s \} : s \in \mathbb{R}_+ \rangle \quad \text{for } i \in N_0,
 \end{aligned}$$

and putting $\rho = \rho^\wedge$, then ρ is a function $N_0 \times \mathbb{R}_+ \rightarrow \mathbb{R}_+$, nondecreasing separately in both arguments, with $\rho(i, 0) = 0$ and $s \leq \rho(i, s)$ for $i \in N_0$ and $s \in \mathbb{R}_+$, and $\rho(i, \cdot)$ continuous at 0, and also such that with

$$B_0 = \sup \{ 1 + |r| : r \in (\chi \cup \partial_2 \chi)[I^{\times 2}] \} \quad \text{we have}$$

$$|(\chi \circ [\text{id}, u] \cdot u \cdot v)^{(i+1)} \cdot s| \leq B_0 (M_0 + 2) M_0 (\mathbf{m} \cdot i^+) + \rho(i, M)$$

for $i \in N_0$ and $s \in I$ and $u, v \in B\mathbf{m}$ and $\mathbf{m} \in (\mathbb{R}^+)^{N_0}$ such that $\mathbf{m} \cdot \emptyset \leq M_0 \leq 1$ and $\sup(\mathbf{m} \cdot i^+) \leq M < +\infty$.

Putting $\theta_0 = \mathbb{R}^+ \times \mathbb{R}_+ \times N_0 \times \mathbb{R}_+ \cap \{(r, s, i, t) :$

$$\rho(i, s) = t(1 - B_0 r(2 + r)) \text{ and } B_0 r(2 + r) < 1 \},$$

we note that $\theta_0(r, \cdot, i)$ is continuous at 0 with $\theta_0(r, 0, i) = 0$ whenever $(r, 0, i) \in \text{dom } \theta_0$. Using this, by a suitable finite induction, one first establishes existence of $\mathbf{n} \in (\mathbb{R}^+)^{l_0+1}$ with $\{(\mathbf{n} \cdot \emptyset, \mathbf{n} \cdot i, i, \mathbf{n} \cdot i^+) : i \in l_0\} \subseteq \theta_0$ and $\mathbf{n} \cdot \emptyset \leq \frac{1}{3} B_0^{-1}$ and $l_0(\mathbf{n} \cdot l_0) \leq 1$, and then fixes one such $\mathbf{n}$. With

$$\mathbf{x}_0 = \langle \sup \{ 1 + |x^{(l)} \cdot s| : s \in I \text{ and } l \in i^+ \} : i \in N_0 \rangle \quad \text{and}$$

$$\mathbf{x}_1 = \langle (\mathbf{n} \cdot \emptyset)^{-1} (\mathbf{x}_0 \cdot l_0)^{-1} (\mathbf{x}_0 \cdot i) : i \in N_0 \rangle \quad \text{and}$$

$$\theta = \{ (r, s, i, t) : \exists t_1 ; (r, s, i, t_1) \in \theta_0 \text{ and } t = \max \{ t_1, \mathbf{x}_1 \cdot i^+ \} \} \quad \text{and}$$

$$\mathcal{M} = (\mathbb{R}^+)^{N_0} \cap \{ \mathbf{m} : \mathbf{n} \subseteq \mathbf{m} \text{ and } \forall i \in N_0 ; \theta(\mathbf{m} \cdot \emptyset, \mathbf{m} \cdot i, i) \leq \mathbf{m} \cdot i^+ \},$$

we now take $\mathcal{B} = \{ B\mathbf{m} : \mathbf{m} \in \mathcal{M} \}$.

Note that by $\mathbf{m} \cdot i \leq \rho(i, \mathbf{m} \cdot i) \leq \theta(\mathbf{m} \cdot \emptyset, \mathbf{m} \cdot i, i) \leq \mathbf{m} \cdot i^+$ every $\mathbf{m} \in \mathcal{M}$ is nondecreasing. By our construction, it is straightforward to verify that (*) above holds. Since every $B \in \mathcal{B}$ is absolutely convex, to have $\mathcal{B}$ a bornological generator for E , one should verify

$$(i_1) \quad \forall \mathbf{m}_1, \mathbf{m}_2 \in \mathcal{M} ; \exists \mathbf{m} \in \mathcal{M} ; \forall i \in N_0 ; \max \{ \mathbf{m}_1 \cdot i, \mathbf{m}_2 \cdot i \} \leq \mathbf{m} \cdot i,$$

$$(i_2) \quad \forall \mathbf{b} \in (\mathbb{R}^+)^{N_0} ; \exists \varepsilon \in \mathbb{R}^+, \mathbf{m} \in \mathcal{M} ; \forall i \in N_0 ; \varepsilon(\mathbf{b} \cdot i) \leq \mathbf{m} \cdot i.$$

To get (i_1) , for $i \in N_0 \setminus l_0$ one applies the recursion

$$\mathbf{m} \cdot i^+ = \max \{ \mathbf{m}_1 \cdot i^+, \mathbf{m}_2 \cdot i^+, \theta(\mathbf{n} \cdot \emptyset, \mathbf{m} \cdot i, i) \} \quad \text{with } \mathbf{m} \cdot l_0 = \mathbf{n} \cdot l_0.$$

For (i_2) one first chooses $\varepsilon \in \mathbb{R}^+$ so that $l_0^+ \subseteq \{i : \varepsilon(\mathbf{b}^i) \leq \mathbf{n}^i\}$, and then applies the recursion $\mathbf{m}^i = \max\{\varepsilon(\mathbf{b}^i), \theta(\mathbf{n}^i, \mathbf{m}^i)\}$. $\square$

To give some perspective, we conclude with the following

11 Remark. Original formulations of Colombeau’s “tameness” conditions are given in [1] and reproduced in [4; Section XIII.4]. There these conditions concern Silva differentiable maps between convex bornological vector spaces. In [9], the conditions are adapted for maps between bornological locally convex spaces. Definition 2.1 in [9; p. 428] introduces certain order k differentiability classes which by [7; Theorem 2.8.1(2), pp. 102, 105] and by a suitable adaptation of the idea in the proof of [8; Theorem 5.20, pp. 62, 27] are precisely the classes $C_c^k(\mathbf{R})$. It seems that in [9] one has not taken into account carefully enough the fact that a Silva C^1 -map need not be continuous with respect to the locally convex topologies when the domain space is not cm-convenient.

References

- [1] J.-F. COLOMBEAU: ‘Fonctions implicites et équations différentielles dans les espaces bornologiques’ *C. R. Acad. Sci. Paris* **272** (1971) 240–243.
- [2] A. FRÖLICHER and A. KRIEGL: *Linear Spaces and Differentiation Theory*, Wiley, Chichester 1988.
- [3] S. HILTUNEN: ‘Differentiation, Implicit functions, and applications to generalized well-posedness’ *preprint*, <http://arXiv.org/abs/math.FA/0504268>.
- [4] H. HOGBE-NLEND: *Théorie des Bornologies et Applications*, Lecture Notes in Math. 273, Springer, Berlin 1971.
- [5] ———: *Bornologies and Functional Analysis*, Math. Studies 26, North Holland, Amsterdam 1977.
- [6] H. JARCHOW: *Locally Convex Spaces*, Teubner, Stuttgart 1981.
- [7] H. H. KELLER: *Differential Calculus in Locally Convex Spaces*, Lecture Notes in Math. 417, Springer, Berlin - Heidelberg - New York 1974.
- [8] A. KRIEGL and P. W. MICHOR: *The Convenient Setting of Global Analysis*, Survey 53, Amer. Math. Soc., Providence 1997.
- [9] J. A. LESLIE: ‘Some integrable subalgebras of the Lie algebras of infinite dimensional Lie groups’ *Trans. Amer. Math. Soc.* **333** (1992) no 1, 423–443.
- [10] W. RUDIN: *Functional Analysis*, Tata McGraw-Hill, New Delhi 1978.

HELSINKI UNIVERSITY OF TECHNOLOGY
 INSTITUTE OF MATHEMATICS, U311
 P.O. BOX 1100
 FIN-02015 HUT
 FINLAND
E-mail address: `shiltune@cc.hut.fi`